



Antifungal Susceptibility of 182 *Fusarium* Species Isolates from 20 European Centers: Comparison between EUCAST and Gradient Concentration Strip Methods

Marion Blaize,^{a,b} Anne-Cecile Normand,^a Sebastien Imbert,^a Abdullah M. S. Al-Hatmi,^{c,d} Erja Chryssanthou,^e Sophie Cassaing,^f Christine Schuttler,^g Lilia Hasseine,^h Caroline Mahinc,ⁱ Damien Costa,^j Christine Bonnal,^k Stéphane Ranque,^l Marc Sautour,^m Elisa Rubio,^{n,o} Laurence Delhaes,^p Arnaud Riat,^q Boualem Sendid,^r Lise Kristensen,^s Marcel Brandenberger,^t Dirk Stubbe,^u Sophie Brun,^v Renaud Piarroux,^{a,w} Arnaud Fekkar^{a,b}

^aAP-HP, Groupe Hospitalier La Pitié-Salpêtrière, Service de Parasitologie Mycologie, Paris, France

^bSorbonne Université, INSERM, CNRS, Centre d'Immunologie et des Maladies Infectieuses, CIMI-Paris, Paris, France

^cCentre of Expertise in Mycology, Radboud University Medical Centre, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands

^dNatural and Medical Sciences Research Center, Department of Microbiology, University of Nizwa, Nizwa, Oman

^eDivision of Clinical Microbiology, Department of Laboratory Medicine, Karolinska Institute, Stockholm, Sweden

^fCHU Toulouse, Service de Parasitologie-Mycologie, Toulouse, France

^gLaboratoire Biogroup-LCD, Levallois, France

^hCHU de Nice, Service de Parasitologie Mycologie, Nice, France

ⁱCHU de Saint-Etienne, Service de Parasitologie Mycologie, Saint-Etienne, France

^jCentre Hospitalier Universitaire de Rouen, Service de Parasitologie Mycologie, Rouen, France

^kAP-HP, Hôpital Bichat-Claude Bernard, Service de Parasitologie Mycologie, Paris, France

^lAix Marseille University, IRD, AP-HM, SSA, VITROME, IHU Méditerranée Infection, Marseille, France

^mCHU de Dijon, Service de Parasitologie Mycologie, Dijon, France

ⁿDepartment of Microbiology, ISGlobal Barcelona Institute for Global Health, Barcelona, Spain

^oCDB, Hospital Clinic, University of Barcelona, Barcelona, Spain

^pCHU de Bordeaux, Groupe Hospitalier Pellegrin, Service de Mycologie, Bordeaux, France

^qGeneva University Hospitals, Bacteriology Laboratory, Geneva, Switzerland

^rCHU de Lille, Department Parasitology-Mycology, Lille, France

^sAarhus University Hospital, Department of Clinical Microbiology, Aarhus, Denmark

^tLabor Synlab Luzern, Lucerne, Switzerland

^uSciensano, BCCM/IHEM Collection, Mycology and Aerobiology Unit, Brussels, Belgium

^vAP-HP, Hôpital Avicenne, Service de Parasitologie-Mycologie, Bobigny, France

^wSorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France

ABSTRACT We determined the susceptibility of 182 *Fusarium* species isolates to five antifungal drugs (amphotericin B, voriconazole, posaconazole, isavuconazole, and terbinafine) by the EUCAST method. Based on the latest taxonomic insights, isolates collected from 20 European centers were distributed into seven complexes and 27 species. The susceptibility was variable, depending on the species. Comparison with the gradient concentration strip method, which was used for 77 isolates, showed essential agreement values for voriconazole, posaconazole, isavuconazole, and amphotericin B of 17%, 91%, 83%, and 70%, respectively.

KEYWORDS *Fusarium*, antifungal susceptibility, comparison, EUCAST, gradient concentration strips, amphotericin B, voriconazole, posaconazole, isavuconazole, terbinafine, Etest

The environmental fungi *Fusarium* spp. are responsible for opportunistic infections in humans, animals, and plants (1–4). *Fusarium* spp. are known to be rather unsusceptible to antifungals, but differences in susceptibility profiles for different drugs have been reported, depending on the species (5). Currently, *Fusarium* spp. include over 300 species, grouped into

Citation Blaize M, Normand A-C, Imbert S, Al-Hatmi AMS, Chryssanthou E, Cassaing S, Schuttler C, Hasseine L, Mahinc C, Costa D, Bonnal C, Ranque S, Sautour M, Rubio E, Delhaes L, Riat A, Sendid B, Kristensen L, Brandenberger M, Stubbe D, Brun S, Piarroux R, Fekkar A. 2021. Antifungal susceptibility of 182 *Fusarium* species isolates from 20 European centers: comparison between EUCAST and gradient concentration strip methods. *Antimicrob Agents Chemother* 65:e01495-21. <https://doi.org/10.1128/AAC.01495-21>.

Copyright © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Arnaud Fekkar, arnaud.fekkar@aphp.fr.

Received 5 August 2021

Returned for modification 19 August 2021

Accepted 13 September 2021

Accepted manuscript posted online

20 September 2021

Published 17 November 2021

23 phylogenetic species complexes (SCs) (6). Given the highly complex classification and rapidly evolving taxonomy, there is a need to update susceptibility determinations according to SCs and species as knowledge evolves and new species are described.

We determined the antifungal susceptibility of 182 *Fusarium* sp. isolates distributed into seven SCs and 27 different phylogenetic species. Isolates were collected from 20 European centers between 1 January 2018 and 31 December 2018 and sent to our institution for identification and antifungal susceptibility testing. Identification was based on a DNA phylogenetic tree-based approach (7). The MICs of amphotericin B, voriconazole, posaconazole, isavuconazole, and terbinafine were determined by using the EUCAST standardized methodology for all isolates and by using gradient concentration strip (GCS) methods (except for terbinafine) for 77 isolates belonging to the most frequent species, i.e., those represented by at least 8 isolates. For EUCAST methodology, drugs concentrations ranged from 0.03 to 16 mg/liter for all tested drugs except posaconazole, which ranged from 0.016 to 8 mg/liter. The MIC values were obtained by visual assessment of the turbidity after 48 h of growth at 37°C and were determined as the lowest concentration with complete inhibition of growth. *Candida parapsilosis* strain ATCC 22019 was used for internal quality control in each EUCAST microplate (even for terbinafine, although there is no defined target value). The GCS method was performed using MIC test strips (Liofilchem, Italy) for isavuconazole and Etest strips (bioMérieux, France) for amphotericin B, voriconazole, and posaconazole. Essential agreement (EA) between methods was considered to be achieved when the MIC values were within the range of 2 dilutions. Correlation between methods was assessed using the Pearson's correlation test.

The comprehensive set of results, showing all MIC values determined by the EUCAST and GCS methods for the different antifungal agents and for all analyzed isolates, is presented in Table S1 in the supplemental material. The main results are presented in Table 1. The majority of isolates were distributed within three SCs, namely, the *Fusarium oxysporum* SC (FOSC), the *Fusarium fujikuroi* SC (FFSC), and the *Fusarium solani* SC (FSSC), reflecting the epidemiology of human fusariosis (7, 8). As expected, *Fusarium* spp. were characterized by low *in vitro* susceptibility to antifungal drugs. However, significant variations in MIC values between the different SCs were observed. Furthermore, within a given SC, some heterogeneity in MIC values was observed, which seemed to be not related to the species identification (see Table S1).

As previously reported, low susceptibility of the FSSC appears to be the rule, regardless of the antifungal drug considered, including the most recently marketed azole isavuconazole (9). Interestingly, isolates of the FFSC were distinguished by greater susceptibility to terbinafine, compared to isolates from other complexes. Overall, amphotericin B was the drug with the lowest geometric mean MIC value (2.3 mg/liter).

EUCAST (readout at 48 h) and GCS (readout at 24, 48, or 72 h) methods showed rather good correlation overall, with the exception of voriconazole. At the 48-h readout, the EA values for voriconazole, posaconazole, isavuconazole, and amphotericin B were 17%, 91%, 83%, and 70%, respectively. However, some variations according to the drug and the incubation period for the GCS method were noticed. For posaconazole and isavuconazole, good correlation was achieved within the first 24 h of growth ($\rho = 0.58$ [$P < 0.0001$] and $\rho = 0.64$ [$P < 0.0001$], respectively); for amphotericin B, the best correlation was obtained after 72 h of growth ($\rho = 0.6$ [$P < 0.0001$]). Lastly, for voriconazole, the correlation coefficients were lower, although statistically significant, whatever the growth period ($\rho = 0.3$ [$P < 0.008$] after 48 h and $\rho = 0.5$ [$P < 0.0001$] after 72 h).

Our study includes a very large number of molecularly identified isolates of *Fusarium* spp. collected in 20 European centers. To the best of our knowledge, these results represent the largest series of *Fusarium* sp. isolates for which a reference method for antifungal susceptibility testing has been used. This work presents a number of pitfalls and limitations. In particular, the interpretation is challenged by the fact that important data, such as categorical endpoints or clinical breakpoints, are still lacking for *Fusarium* spp. Espinel-Ingroff et al. proposed epidemiological cutoff values (ECVs) based on CLSI methods for two SCs plus the species *Fusarium verticilloides* (10), but there is still a long way to go. In addition, some species were represented by a small number of isolates, making it impossible to produce reliable aggregate data. Finally,

TABLE 1 MIC values determined by EUCAST and GCS methods for 182 clinical isolates of *Fusarium* spp.

Method, SC, and drug	MIC range (mg/liter)	MIC geometric mean (mg/liter)	No. of isolates with MIC of ^a :								EA (%) vs EUCAST method ^b			
			0.125 mg/liter	0.25 mg/liter	0.5 mg/liter	1 mg/liter	2 mg/liter	4 mg/liter	8 mg/liter	16 mg/liter ^c	> 16 mg/liter	method ^b		
EUCAST method ^d														
Total (182 isolates)														
TBF	1 to >16	13.8	0	0	0	3	17	38	12	0	112			
VCZ	2 to >16	14.3	0	0	0	0	6	22	43	35	76			
PSZ	0.5 to >8	13.1	0	0	3	5	6	0	0	168				
ICZ	2 to >16	24.6	0	0	0	0	2	3	12	28	137			
AMB	0.25 to >16	2.3	0	1	5	62	61	16	19	15	3			
FFSC (54 isolates)														
TBF	1 to >16	3.9	0	0	0	2	13	31	4	0	4			
VCZ	2 to >16	11.6	0	0	0	0	2	6	16	21	9			
PSZ	0.5 to >8	11.0	0	0	3	2	2	0	0	47				
ICZ	2 to >16	24.1	0	0	0	0	1	2	4	4	43			
AMB	0.5 to >16	2.6	0	0	2	17	16	4	7	8	0			
FOSC (65 isolates)														
TBF	1 to >16	20.9	0	0	0	1	1	5	8	0	50			
VCZ	2 to >16	11.0	0	0	0	0	2	14	22	6	21			
PSZ	1 to >8	13.9	0	0	0	1	3	0	0	61				
ICZ	2 to >16	20.9	0	0	0	0	1	0	7	22	35			
AMB	0.5 to >16	2.2	0	0	1	21	27	5	6	3	2			
FSSC (53 isolates)														
TBF	>16	32.0	0	0	0	0	0	0	0	0	53			
VCZ	8 to >16	28.4	0	0	0	0	0	0	1	7	45			
PSZ	>8	16.0	0	0	0	0	0	0	0	53				
ICZ	>16	32.0	0	0	0	0	0	0	0	0	53			
AMB	0.25 to >16	1.9	0	1	2	21	15	6	5	2	1			
GCS method ^e														
Total (77 isolates)														
VCZ	0.25 to >16	2.2	0	2	11	15	19	18	4	2	6	17		
PSZ	0.125 to >8	10.8	1	0	7	0	0	1	0	68		91		
ICZ	0.125 to >16	17.2	1	0	0	6	7	1	0	0	62	83		
AMB	0.25 to >16	1.5	0	3	12	24	22	8	6	1	1	70		
FFSC (18 isolates)														
VCZ	0.25 to >16	1.3	0	1	5	4	4	3	0	1	0	22		
PSZ	0.125 to >8	3.8	1	0	6	0	0	0	0	11		72		
ICZ	1 to >16	9.0	0	0	0	5	2	0	0	0	11	72		
AMB	0.5 to >16	2.1	0	0	1	5	8	1	2	1	0	67		
FOSC (20 isolates)														
VCZ	0.25 to >8	0.9	0	1	6	9	3	0	1	0	0	5		
PSZ	0.5 to >8	12.6	0	0	1	0	0	1	0	18		90		
ICZ	0.125 to >16	9.2	1	0	0	1	5	1	0	0	12	60		
AMB	0.5 to >16	2.1	0	0	3	5	5	3	3	0	1	70		

(Continued on next page)

(Continued on next page)

TABLE 1 (Continued)

Method, SC, and drug	MIC range (mg/liter)	MIC geometric mean (mg/liter)	No. of isolates with MIC of ^a :								EA (%) vs EUCAST method ^b	
			0.125 mg/liter	0.25 mg/liter	0.5 mg/liter	1 mg/liter	2 mg/liter	4 mg/liter	8 mg/liter	16 mg/liter ^c	>16 mg/liter	
FSSC (39 isolates)												
VCZ	1 to >16	4.5	0	0	0	2	12	15	3	1	6	21
PSZ	>8 to >8	16.0	0	0	0	0	0	0	0	39		100
ICZ	>16 to >16	32.0	0	0	0	0	0	0	0	0	39	100
AMB	0.25 to 8	1.1	0	3	8	14	9	4	1	0	0	72

^aMIC values for the GCS method were obtained with a 48-h readout. TBF, terbinafine; AMB, amphotericin B; ICZ, isavuconazole; PSZ, posaconazole; VCZ, voriconazole.

^bEA between methods was considered to be achieved when the MIC values were within ± 2 dilutions.

^cFor posaconazole, concentrations ranged from 0.016 to 8 mg/liter; therefore, this column should be interpreted as ≥ 8 mg/liter.

^dTotal, FFSC, FOSC, FSSC, *Fusarium culmorum* SC, *Fusarium dimerum* SC, *Fusarium incarnatum* SC, and *Fusarium redolens* SC; FFSC, *Fusarium verticillioides*, *Fusarium proliferatum*, *Fusarium sacchari*, *Fusarium acutatum*, *Fusarium andiyazi*, and *Fusarium lactis*; FOSC, *Fusarium nirenbergiae*, *Fusarium curvatum*, *Fusarium elaeidis*, *Fusarium gossypinum*, *Fusarium languescens*, and *Fusarium triseptatum*; FSSC, *Fusarium falciforme*, *Fusarium keratoplasticum*, *Fusarium petrophilum*, FSSC new species 1, FSSC new species 2, *Fusarium lichenichola*, *Fusarium robiniae*, *F. solani* FSSC5, and *F. solani* FSSC9; FOSC, *Fusarium oxysporum* species complex; FFSC, *Fusarium solani* species complex.

^eTotal, FFSC, FOSC, and FSSC; FFSC, *F. verticillioides* and *F. proliferatum*; FOSC, *F. nirenbergiae*; FSSC, *F. falciforme*, *F. keratoplasticum*, *F. petrophilum*, FSSC new species 1, and *F. solani* FSSC5.

whether clinical breakpoints will be available some day is uncertain, as a clear and definite relationship between MICs obtained *in vitro* and therapeutic efficacy during the course of fusariosis is doubtful (11, 12). Similarly, a recent study found no correlation between MIC values and mortality rates at day 90 (4) (although this point does not take into consideration the given treatment), while another report indicated that a patient with fusariosis due to an isolate with a high itraconazole MIC showed clinical improvement after treatment with this antifungal agent (13). In any case, beyond the potential clinical impact, the importance of conducting epidemiological surveys of *Fusarium* sp. susceptibility to antifungal agents has two additional justifications. The first is the rapid evolution of taxonomy and the fact that data generated at one time may become inaccurate soon thereafter. The second is the environmental nature of *Fusarium* spp., because of which they are prone to have their resistance profiles evolve according to their exposure to antifungal agents used in agriculture (especially demethylase inhibitors), as is the case with *Aspergillus fumigatus*.

In a previous work, EUCAST and GCS methods showed an EA above 85% but the study included only 20 *Fusarium* sp. isolates and as many different species (14). For us, the results of the EUCAST and GCS methods were correlated. However, similar to what we reported concerning *Aspergillus* section *Nigri* (15), we found consistently lower MIC values with the GCS method, compared with the EUCAST method, particularly for azole drugs. Consequently, the EA values were variable, being particularly low for voriconazole (<25%) and more suitable (70% to 91%) for the other azole drugs. As our work provides new data on the susceptibility of *Fusarium* spp. to antifungal agents, we hope that it can contribute in the future to the establishment of accurate ECVs for a wider range of species.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, XLSX file, 0.03 MB.

REFERENCES

- Gleason FH, Allerstorfer M, Lilje O. 2020. Newly emerging diseases of marine turtles, especially sea turtle egg fusariosis (SEFT), caused by species in the *Fusarium solani* complex (FSSC). *Mycology* 11:184–194. <https://doi.org/10.1080/21501203.2019.1710303>.
- Triest D, Hendrickx M. 2016. Postharvest disease of banana caused by *Fusarium musae*: a public health concern? *PLoS Pathog* 12:e1005940. <https://doi.org/10.1371/journal.ppat.1005940>.
- Poignon C, Blaize M, Vezinet C, Lampros A, Monsel A, Fekkar A. 2020. Invasive pulmonary fusariosis in an immunocompetent critically ill patient with severe COVID-19. *Clin Microbiol Infect* 26:1582–1584. <https://doi.org/10.1016/j.cmi.2020.06.026>.
- Nucci M, Jenks J, Thompson GR, Hoenigl M, dos Santos MC, Forghieri F, Rico JC, Bonuomo V, López-Soria L, Lass-Flörl C, Candoni A, García-Vidal C, Cattaneo C, Buil J, Rabagliati R, Roiz MP, Gudiol C, Fracchiolla N, Campos-Herrero MI, Delia M, Farina F, Fortun J, Nadali G, Sastre E, Colombo AL, Pérez Nadeles E, Alastruey-Izquierdo A, Pagano L. 2021. Do high MICs predict the outcome in invasive fusariosis? *J Antimicrob Chemother* 76:1063–1069. <https://doi.org/10.1093/jac/dkaa516>.
- Al-Hatmi AMS, Bonifaz A, Ranque S, Sybren de Hoog G, Verweij PE, Meis JF. 2018. Current antifungal treatment of fusariosis. *Int J Antimicrob Agents* 51:326–332. <https://doi.org/10.1016/j.ijantimicag.2017.06.017>.
- Geiser DM, Al-Hatmi AMS, Aoki T, Arie T, Balmas V, Barnes I, Bergstrom GC, Bhattacharyya MK, Blomquist CL, Bowden RL, Brankovics B, Brown DW, Burgess LW, Bushley K, Busman M, Cano-Lira JF, Carrillo JD, Chang HX, Chen CY, Chen W, Chilvers M, Chulze S, Coleman JJ, Cuomo CA, de Beer ZW, de Hoog GS, Del Castillo-Múnera J, Del Ponte EM, Diéguez-Urbeondo J, Di Pietro A, Edel-Hermann V, Elmer WH, Epstein L, Eskalen A, Esposto MC, Everts KL, Fernández-Pavía SP, da Silva GF, Foroud NA, Fourie G, Frandsen RJN, Freeman S, Freitag M, Frenkel O, Fuller KK, Gagkaeva T, Gardiner DM, Glenn AE, Gold SE, Gordon TR, Gregory NF, Gryzenhout M, Guarro J, Gugino BK, Gutierrez S, Hammond-Kosack KE, Harris LJ, Homa M, Hong CF, Hornok L, Huang JW, Ilkit M, Jacobs A, Jacobs K, Jiang C, Jiménez-Gasco MDM, Kang S, Kasson MT, Kazan K, Kennell JC, Kim HS, Kistler HC, Kuldau GA, Kulik T, Kurzai O, Laraba I, Laurence MH, Lee T, Lee YW, Lee YH, Leslie JF, Liew ECY, Lofton LW, Logrieco AF, López-Berges MS, Luque AG, Lysøe E, Ma LJ, Marra RE, Martin FN, May SR, McCormick SP, McGee C, Meis JF, Migheli Q, Nor NMIM, Monod M, Moretti A, Mostert D, Mulé G, Munaut F, Munkvold GP, Nicholson P, Nucci M, O'Donnell K, Pasquali M, Pfenning LH, Prigitano A, Proctor RH, Ranque S, Rehner SA, Rep M, Rodríguez-Alvarado G, Rose LJ, Roth MG, Ruiz-Roldán C, Saleh AA, Salleh B, Sang H, Scandiani MM, Scauflaire J, Schmale DG, III, Short DPG, Šišić A, Smith JA, Smyth CW, Son H, Spahr E, Stajich JE, Steenkamp E, Steinberg C, Subramaniam R, Suga H, Summerell BA, Susca A, Swett CL, Toomajian C, Torres-Cruz TJ, Tortorano AM, Urban M, Vaillancourt LJ, Vallad GE, van der Lee TAJ, Vanderpool D, van Diepeningen AD, Vaughan MM, Venter E, Vermeulen M, Verweij PE, Viljoen A, Waalwijk C, Wallace EC, Walther G, Wang J, Ward TJ, Wickes BL, Wiederhold NP, Wingfield MJ, Wood AKM, Xu JR, Yang XB, Yli-Mattila T, Yun SH, Zakaria L, Zhang H, Zhang N, Zhang SX, Zhang X. 2021. Phylogenomic analysis of a 55.1 kb 19-gene dataset resolves a monophyletic *Fusarium* that includes the *Fusarium solani* species complex. *Phytopathology* PHYTO08200330LE. <https://doi.org/10.1094/PHYTO-08-20-0330-LE>.
- Normand A-C, Imbert S, Brun S, Al-Hatmi AMS, Chrysanthou E, Cassaing S, Schuttler C, Hasseine L, Mahinc C, Costa D, Bonnal C, Ranque S, Sautour M, Rubio E, Delhaes L, Riat A, Sendid B, Kristensen L, Brandenberger M, Guitard J, Packeu A, Piarroux R, Fekkar A. 2021. Clinical origin and species distribution of *Fusarium* spp. isolates identified by molecular sequencing and mass spectrometry: a European multicenter hospital prospective study. *J Fungi (Basel)* 7:246. <https://doi.org/10.3390/jof7040246>.
- Pérez-Nadeles E, Alastruey-Izquierdo A, Linares-Sicilia MJ, Soto-Debrán JC, Abdala E, García-Rodríguez J, Montejo M, Muñoz P, Lletí MS, Rezusta A, de Pipaón MRP, Yáñez L, Merino E, Campos-Herrero MI, Costa-Mateo JM, Fortún J, García-Lozano T, García-Vidal C, Fernández-Ruiz M, Sánchez-Reus F, Castro-Méndez C, Guerrero-Lozano I, Soler-Palacín P, Aguado JM, Martínez-Martínez L, Torre-Cisneros J, Nucci M. 2021. Invasive fusariosis in nonneutropenic patients, Spain, 2000–2015. *Emerg Infect Dis* 27:26–35. <https://doi.org/10.3201/eid2701.190782>.

9. Herkert PF, Al-Hatmi AMS, de Oliveira Salvador GL, Muro MD, Pinheiro RL, Nucci M, Queiroz-Telles F, de Hoog GS, Meis JF. 2019. Molecular characterization and antifungal susceptibility of clinical *Fusarium* species from Brazil. *Front Microbiol* 10:737. <https://doi.org/10.3389/fmicb.2019.00737>.
10. Espinel-Ingroff A, Colombo AL, Cordoba S, Dufresne PJ, Fuller J, Ghannoum M, Gonzalez GM, Guarro J, Kidd SE, Meis JF, Melhem TMS, Pelaez T, Pfaller MA, Szeszs MW, Takahaschi JP, Tortorano AM, Wiederhold NP, Turnidge J. 2016. International evaluation of MIC distributions and epidemiological cutoff value (ECV) definitions for *Fusarium* species identified by molecular methods for the CLSI broth microdilution method. *Antimicrob Agents Chemother* 60:1079–1084. <https://doi.org/10.1128/AAC.02456-15>.
11. Navarro-Rodríguez P, Guevara-Suarez M, Paredes K, Celis A, Guarro J, Capilla J. 2018. Lack of correlation of ECV and outcome in an in vivo murine model of systemic fusariosis. *Diagn Microbiol Infect Dis* 92:124–126. <https://doi.org/10.1016/j.diagmicrobio.2018.05.019>.
12. Al-Hatmi AMS, Curfs-Breuker I, de Hoog GS, Meis JF, Verweij PE. 2017. Antifungal susceptibility testing of *Fusarium*: a practical approach. *J Fungi (Basel)* 3:19. <https://doi.org/10.3390/jof3020019>.
13. Al-Hatmi AMS, Bonifaz A, Tirado-Sánchez A, Meis JF, de Hoog GS, Ahmed SA. 2017. *Fusarium* species causing eumycetoma: report of two cases and comprehensive review of the literature. *Mycoses* 60:204–212. <https://doi.org/10.1111/myc.12590>.
14. Al-Hatmi AMS, Normand A-C, Ranque S, Piarroux R, de Hoog GS, Meletiadis J, Meis JF. 2017. Comparative evaluation of Etest, EUCAST, and CLSI methods for amphotericin B, voriconazole, and posaconazole against clinically relevant *Fusarium* species. *Antimicrob Agents Chemother* 61:e01671-16. <https://doi.org/10.1128/AAC.01671-16>.
15. Carrara B, Richards R, Imbert S, Morio F, Sasso M, Zahr N, Normand AC, Pape PL, Lachaud L, Ranque S, Maubon D, Piarroux R, Fekkar A. 2020. Species distribution and comparison between EUCAST and gradient concentration strips methods for antifungal susceptibility testing of 112 *Aspergillus* section *Nigri* isolates. *Antimicrob Agents Chemother* 64:e02510-19. <https://doi.org/10.1128/AAC.02510-19>.